DATE ORDERED	TIME ORDERED	PHYSICIAN'S ORDER
		DVT Prophylaxis Protocol
		□ No Risk Assessed □ Patient on Anticoagulants
- 62		Low Risk:
		□ Early ambulation ad lib
s		□ TEDS, knee high
		□ TEDS, thigh high
		□ VENODYNES
		Moderate Risk:
		□ Venodynes until patient ambulates without assistance
		☐ Heparin 5000 units sq q 12h x days
		□ Lovenox 30 mg sc q 12h, first dose to begin hours post op
		(Non-Frail Arthroplasty patients only – Hold if active surgical site bleeding, call MD)
		□ Lovenox 40 mg qd, first dose to begin hours post op (Non-Frail
		Arthroplasty patients only – Hold if active surgical site bleeding, call MD)
		□ Other
		High Risk:
		□ Venodynes until patient ambulates without assistance
		One of the following if anticoagulation not otherwise contraindicated.
		Coumadin protocol for prophylaxis (refer to coumadin protocol order sheet protocol)
		□ Non-thrombolytic Heparin Protocol order sheet.
		□ Lovenox 30 mg sc q 12h, first dose to begin hours post op
		(Non-Frail Arthroplasty patients only – Hold if active surgical site bleeding, call MD)
		□ Lovenox 40 mg sc qd, first dose to begin hours post op (Non-Frail
		Arthroplasty patients only – Hold if active surgical site bleeding, call MD)
		□ Radiology consult for Inferior vena caval filter (Multi-trauma and high risk
		Patient with recent DVT/PE only – and if anticoagulation contraindicated).
		☐ Heparin 5000 units sq q 12h x days (elderly frail hip fracture patients)
		□ Other
		MD Signature Date
		Must complete Risk Assessment on back over
NSTRUC	TIONS: 1.1	MPRINT PATIENT'S PLATE IN DESIGNATED AREA BEFORE PLACING IN CHART ADDRESSOGRAPH
-		